

Expert Opinion

1. Introduction
2. Effect of NHE-1 on cellular level
3. NHE-1 and vascular risk factors
4. Clinical trials and potential therapeutic applications
5. Modulation of NHE-1 activity by clinically used compounds
6. Expert opinion

Na⁺/H⁺ exchanger-1: a link with atherogenesis?

Maria Sarigianni, Apostolos Tsapas, Dimitri P Mikhailidis, Martha Kaloyianni, George Koliakos, Larry Fliegel & Konstantinos Paletas[†]

[†]*Aristotle University of Thessaloniki, Medical School, Metabolic Diseases Unit, Second Department of Internal Medicine, 15 Ag. Sofias str, 54623, Thessaloniki, Greece*

Importance of the field: The sodium/hydrogen exchanger-1 (NHE-1/SLC9A1) is a ubiquitous plasma membrane protein whose main role is maintenance of intracellular pH and volume. NHE-1 plays a role in atherogenesis; however, its clinical relevance has not yet been established.

Areas covered in this review: We herein review the contribution of NHE-1 in atherogenesis (namely its effect on endothelial cells, monocytes, smooth muscle cells and platelets).

What the reader will gain: Studies have shown that NHE is involved in atherogenesis-related properties of isolated monocytes. We also consider the relationship between NHE-1 and vascular risk factors such as obesity, diabetes mellitus, hypertension, dyslipidemia and inflammation.

Take home message: Even though clinical trials with certain NHE-1 inhibitors have had discouraging results, NHE-1 cannot be excluded as a potential future therapeutic target for the prevention and/or treatment of atherosclerosis.

Keywords: atherosclerosis, cariporide, endothelial cells, monocytes, platelets, smooth muscle cells, sodium-hydrogen antiporter

Expert Opin. Investig. Drugs (2010) 19(12):1545-1556

1. Introduction

The sodium/hydrogen exchanger-1 (NHE-1/SLC9A1) is a ubiquitous integral membrane protein expressed in mammalian cells [1]. Its main role is intracellular pH maintenance, which is achieved by exchanging 1 intracellular H⁺ for 1 extracellular Na⁺ [1,2]. It is also important for cell volume maintenance and cytoskeletal reorganization. Furthermore, it takes part in cell proliferation, apoptosis and migration [1]. NHE-1 activity is stimulated by intracellular acidosis, hormones and growth factors and inhibited by amiloride derivatives [3].

NHE-1 plays a role in atherogenesis; however, its clinical relevance has not yet been established. Atherogenesis is an inflammatory process starting at birth, which progresses continuously as age increases. However, the presence of risk factors such as obesity, diabetes, dyslipidemia, hypertension and smoking accelerates this process. Atherogenesis begins with endothelial dysfunction which attracts and promotes monocyte adhesion to the endothelium. Monocytes transmigrate through the endothelium in the sub-endothelial space, differentiate into macrophages, phagocytose oxidized low-density lipoprotein (oxLDL) and become foam cells. Simultaneously, smooth muscle cells (SMCs) migrate in the sub-endothelial space, phagocytose the oxLDL and transform also into foam cells. Atheromatous plaque is formed with a fibrous cap. Fibrous plaques are often vulnerable to rupture causing thrombosis and stenosis [4-6].

We herein review the contribution of NHE-1 in atherogenesis (namely its effect on endothelial cells, monocytes, SMCs and platelets). We also consider the relationship between NHE-1 and vascular risk factors such as obesity, diabetes mellitus, hypertension, dyslipidemia and inflammation. Even though clinical trials with

informa
healthcare

Article highlights.

- The sodium/hydrogen exchanger-1 (NHE-1) is a ubiquitous plasma membrane protein whose main role is to maintain intracellular pH and volume.
- NHE-1 plays a role in monocyte differentiation, adhesion, migration, CD36 expression and phagocytosis of oxidized low-density lipoprotein.
- NHE-1 can influence smooth muscle cell and platelet function.
- NHE-1 activity is increased in red blood cells obtained from obese humans.
- High concentrations of glucose or insulin can activate NHE-1 suggesting that NHE-1 activity might be increased in diabetes.
- Whether NHE-1 inhibition can be used to treat hypertension remains to be established.
- Cholesterol can influence NHE-1 activity.
- NHE-1 inhibition can improve inflammatory markers.
- Several commonly used drugs (e.g., enalapril, nifedipine, lisinopril and valsartan) seem to modulate NHE-1 activity.
- Even though clinical trials with NHE-1 inhibitors reported discouraging results in terms of vascular events, these drugs merit further investigation.

This box summarizes key points contained in the article.

certain NHE-1 inhibitors had discouraging results in the treatment of ischemic cardiac events, NHE-1 cannot be excluded as a potential future therapeutic target [7-10]. It has to be noted that there is no known clinical trial for the evaluation of NHE1 inhibition for the prevention or treatment of atherosclerosis.

2. Effect of NHE-1 on cellular level

2.1 Endothelial cells

Endothelial cells form the inner lining of the vasculature [4]. Circulating leukocytes normally do not adhere to the endothelium, mainly due to local production of NO, prostacyclin and CD39 [11]. Dysfunctional endothelium promotes monocyte rolling, adherence and transmigration into the sub-endothelial space through increased expression of adhesion molecules (e.g., intracellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule, P- and E-selectins and integrins) and chemokines (e.g., monocyte chemoattractant protein-1, MCP-1) [4].

Endothelial function is influenced by NHE-1 activity. Inhibition of NHE-1 preserves endothelial-dependent vasorelaxation after ischemia and prevents endothelial dysfunction. When reperfused after ischemia, isolated rat hearts showed impaired endothelial-dependent acetylcholine-induced vasorelaxation, which was prevented with SM-20550 (a NHE-1 inhibitor) administration pre- and post-ischemia [12]. Similar findings have been reported in animal models with other NHE-1 inhibitors such as cariporide [13-15],

benzamide, N-(aminoiminomethyl)-4-[4-(2-furanylcarbonyl)-1-piperazinyl]-3-(methylsulfonyl), methanesulfonate] (BIIB-513) [16] and eniporide [17]. In the last study, eniporide preserved endothelial function (estimated by means of acetylcholine-induced vasorelaxation) after 90 min of ischemia in dog hearts. NHE inhibition in *ex vivo* cultures protected endothelial cells against cell injury and death [12]; thus, it could prove beneficial in states of endothelial dysfunction.

In contrast with these findings, another study reported that *in vivo* inhibition of NHE in rat basilar artery prevented the acetylcholine- and bradykinin-induced vasodilatation but did not affect the baseline artery diameter. Hence, intracellular alkalization mediated by NHE activation could enable NO production in basilar artery endothelium [18]. Yet, the reason for this paradoxical finding is not obvious. It could reflect response variability in different vascular beds, which could be attributed to variations in receptor distribution according to the anatomical location of arteries in experimental models [19]. Another possible explanation is the pre-existing ischemia in some experiments which predispose to a different role of NHE. Ischemia-reperfusion promotes the local production of free radicals (reactive oxygen species (ROS)) and an interaction between NHE1-related signaling and ROS has been reported [20,21]. Heat shock protein 70, a molecule related to oxidative stress, seems to be attached to the regulatory domain of NHE; however, the role of this association is not yet clear [22].

Endothelial dysfunction in subjects with diabetes might also be related to NHE [23]. High glucose concentrations activate NHE, thereby, inhibiting NO production via a dissociation of eNOS from heat shock protein 90 in HUVECs [23]. Cariporide (a specific inhibitor of NHE-1) [24] and siRNA for NHE reversed the above-mentioned results [23]. Moreover, administration of cariporide in diabetic animal models has been associated with coronary endothelium protection [25].

The endothelium produces a number of vasoactive substances (e.g., NO, prostacyclin) and NHE-1 could play a role in this process [26]. However, inhibition of NHE does not affect basal production of platelet derived growth factor (PDGF) and monocyte adhesion to endothelial cells but does inhibit thrombin-induced PDGF production by human aortic endothelial cells [27].

The interaction between endothelial cells and leukocytes (mainly neutrophils) is influenced by NHE-1. Cariporide inhibits leukocyte rolling, adhesion and transmigration in states such as post-ischemia reperfusion, where these leukocyte properties are enhanced. However, it has no effect in normal states where leukocyte rolling, adhesion and transmigration are not increased [28]. Thus, pharmacological inhibition is beneficial only when it is highly selective in disease states where undesirable effect occurs (e.g., excessive leukocyte transmigration). The aforementioned interaction between endothelium and leukocytes is regulated by the expression of adhesion molecules such as ICAM-1, P- and E-selectins as

well as MCP-1. Inhibition of NHE-1 counteracts the high glucose-induced increase in ICAM-1 expression in endothelial cells and the subsequent monocyte adhesion to endothelial cells [29]. P-selectin expression is also inhibited by cariporide in states of increased expression [28]. Similar findings have been reported for E-selectin and MCP-1 when NHE-1 is inhibited by amiloride [30]. Other blood cells such as platelets are also influenced by NHE-1 inhibition and show decreased adhesion to endothelium following cariporide administration [31].

Furthermore, amiloride derivatives, mainly 5-(*N,N*-hexamethylene)-amiloride, cause apoptosis of endothelial cells and upregulation of COX-2 (which is involved in the inflammatory response of endothelial cells) and several genes in endoplasmic reticulum [32].

2.2 Monocytes

Monocytes play a key role in atherogenesis following their recruitment and attachment to the activated endothelial cells through increased expression of leukocyte adhesion molecules [5,33,34]. Subsequently, monocytes migrate into the intima media under the influence of chemokines, differentiate to macrophages, proliferate and form foam cells [4,5].

NHE plays a significant role in the differentiation process of monocytes. Studies with HL60 promyelocytic leukemia cell lines which differentiate into monocytes/macrophages have shown that cell differentiation is preceded by a NHE mRNA and protein synthesis increase [35]. Similarly, in stem cells which were treated with dimethyl sulfoxide in order to differentiate to cardiomyocytes, NHE-1 inhibition prevented cell differentiation [36] and increased expression of NHE-1 from an adenoviral vector promoted cardiomyocyte cell differentiation [37]. Moreover, several monocyte properties involved in atherogenesis seem to be mediated by NHE-1. These have all been examined following enhancement by a hormone. NHE-1 inhibition has been shown to prevent this enhancement. Leptin, adrenaline, endothelin-1 and high glucose concentrations increased monocyte adhesion, migration, CD36 expression and phagocytosis of oxLDL via NHE-1-dependent mechanisms [21,38-42]. Furthermore, inhibition of NHE-1 activity with cariporide prevents the lysophosphatidylcholine (LPC)-induced adhesion of monocytes to bovine aortic SMCs [43]. It is worth noting that cariporide has no effect on monocyte adhesion to SMCs in the absence of LPC [43]. This finding is similar to the one reported for endothelial cells where NHE-1 inhibition seems to have a beneficial effect only when there is activation of a process but not under basal conditions.

2.3 SMCs

SMCs influence plaque stability [44]. SMCs under the influence of growth factors migrate into the intima [45] and change their phenotype from contractile to secretory. These secretory SMCs proliferate and produce collagen and several inflammatory mediators enhancing the atherogenic process [4,46].

NHE-1 has a regulatory role in the maintenance of intracellular pH (pHi) in SMCs [47]. The physiologic function of the vascular wall is influenced by SMC pHi; acute intracellular acidification causes contraction [48,49]. Activation of NHE-1 in SMCs has been observed following *in vitro* addition of noradrenaline and phenylephrine suggesting an involvement of pHi in SMC contraction [50]. Furthermore, SMC growth is modulated by pHi [51]. NHE is localized in the plasma membrane but is also found at the nuclear membrane [52]; thus, it could play a role in the function of nucleus in SMCs.

Studies report that NHE-1 mediates several aspects of SMC function such as proliferation, adhesion, migration, molecule expression and apoptosis. Inhibition of NHE-1 with a selective inhibitor, sabiporide, results in the arrest of cell cycle progression at the G₀/G₁ phase and inhibits pulmonary SMC proliferation. Under basal conditions, SMCs are in non-proliferative state while in atherosclerosis they proliferate. Arresting the cell cycle of SMCs could prevent the progression of atherogenesis. In the same cell type, sabiporide inhibits cell migration in a dose-dependent manner [53] indicating that NHE-1 is involved in SMC migration. The level of NHE-1 message varies in response to the state of SMC proliferation. NHE-1 mRNA are increased in SMCs during exponential growth, but decrease at confluency. SMCs deficient in NHE-1 have reduced growth. Injury to the vascular wall increases NHE-1 mRNA levels as does pressure overload of hearts. These results suggest that NHE-1 is involved in vascular SMC proliferation [54].

NHE-1 seems to also be involved in LPC-induced increase in ICAM-1 expression by SMCs and monocyte adhesion to them [43] as cariporide has been shown to inhibit ICAM-1 expression on SMCs in a dose-dependent manner [43]. However, ICAM-1 expression on SMCs is not affected by cariporide in the absence of LPC [43].

NHE-1 inhibition causes apoptosis of SMCs [55] possibly due to cellular acidification, a finding that has been shown both in peripheral blood monocytes and leukemic cells [56,57].

There is evidence supporting that NHE-1 could contribute to the differences in vascular risk between males and females. A study reports that low physiologic concentrations of 17- β -estradiol activate NHE-1 and increase pHi in rat SMCs. However, high levels of 17- β -estradiol did not affect pHi which could contribute to the different risk for atherosclerosis in men and postmenopausal women compared with premenopausal women [50]. The activation of NHE-1 is possibly through a nongenomic mechanism in these cells.

2.4 Platelets

Platelets are activated after contact with sub-endothelial layers which contain collagen and extracellular matrix proteins leading to platelet aggregation. Another important role of platelets is the production of inflammatory proteins, chemokines and proliferative factors [4,58,59]. A variety of agonists stimulate NHE activity in platelets including thrombin, arachidonic

acid, platelet activating factor, angiotensin II, endothelin and phorbol esters [60]. Several novel NHE-1 inhibitors have been reported to prevent platelet swelling [61]. An increase in median platelet volume (MPV) is an early phase of platelet activation [62]. Whether NHE-1 inhibitors block agonist-induced increase in MPV remains to be established. Furthermore, it is not known if the platelet swelling that is influenced by NHE-1 inhibitors reflects the same process as an agonist-induced increase in MPV.

Following activation, platelets release microvesicles with pro-coagulant activity [63]. Inhibition of NHE-1 with 5-(*N*-ethyl-*N*-isopropyl) amiloride (EIPA) prevents collagen- and phorbol ester-evoked vesiculation [64]. Stimulation of porcine platelets by thrombin not only results in NHE-1 activation but also in secretion of serotonin [65]. Inhibition of NHE-1 prevents thrombin-induced secretion of serotonin [65,66]. Serotonin is an activator of platelets, a vasoconstrictor and acts as a proliferator of SMCs. Therefore, this bioamine may play a role in atherogenesis and thrombogenesis [67,68].

After platelet activation, phosphatidyl serine (PS) on the plasma membrane is exposed and initiates the coagulation cascade. NHE-1 seems to be involved in this process. Inhibition of NHE-1 with EIPA results in decreased exposure of PS in human platelets [69]. A similar finding has been observed when PS expression was stimulated by arginine vasopressin [70].

Another inhibitor of NHE-1 (KR-32560) prevents collagen-induced platelet aggregation possibly through inhibition of cytoplasmic Ca²⁺ mobilization and arachidonic acid liberation [71]. Similarly, inhibition of NHE-1 with EIPA prevents desmopressin-evoked platelet pro-coagulant activity [72]. Besides platelet aggregation, inhibition of NHE-1 has been shown to decrease platelet-leukocyte aggregation [73].

All these findings indicate that NHE-1 is involved in platelet activation, serotonin release, platelet aggregation and interaction with other cell types. Furthermore, inhibition of NHE-1 seems to favorably influence platelet function suggesting that NHE-1 inhibitors could contribute to atherogenesis prevention and treatment. It appears that the presence of sodium ion in the medium is required for platelet aggregation [60].

3. NHE-1 and vascular risk factors

3.1 Obesity

NHE-1 activity is increased in red blood cells (RBCs) obtained from obese humans and correlates with body mass index [74]. Furthermore, NHE-1 mediates the atherosclerotic properties of monocytes (adhesion, migration, CD36 expression and oxLDL phagocytosis) from obese and lean subjects [39,41,42,75]. A possible link between obesity and accelerated atherogenesis could be the increased NHE-1 activity in this state. This may be attributed to the several hormonal changes associated with obesity, such as increased insulin and leptin plasma levels [76] or increased

sympathetic activity [77-79]. In this regard, efficacy of several hormones in stimulation of NHE-1 activity has been examined, including insulin, leptin and adrenaline. High concentrations of insulin increased NHE-1 activity in human RBCs [80] obtained from lean healthy subjects but not from obese subjects [81]. However, insulin activated NHE-1 in human monocytes obtained from obese subjects [82]. Leptin increased NHE-1 activity in RBCs obtained from lean and obese humans. However, the increase was greater in the lean group compared with the obese group [83]. This difference could potentially be explained by the existence of leptin resistance in RBCs obtained from obese humans. Finally, adrenaline activated NHE-1 in human RBCs and the response was greater in the RBCs obtained from the obese compared with the lean subjects [84].

Though the aforementioned hormones exhibit a different NHE-1 response in obese subjects compared with the lean, the effect varied with adrenaline having a greater stimulatory effect and insulin and leptin having less or no effect on NHE-1 activity. In the obese, insulin, leptin and adrenaline levels may be increased [76,79] which could contribute to altered NHE-1 activity. Interestingly, there are no data whether obese subjects have increased NHE-1 mRNA or protein expression.

3.2 Diabetes

Diabetes is a risk factor for atherosclerosis. High concentrations of glucose activate NHE-1 [40,85] in a time-dependent manner and this effect was inhibited by cariporide [23]. Furthermore, high glucose concentrations induce impairment of endothelium-dependent vasorelaxation in rat aortic rings. This effect is concentration-dependent [86] and attenuates after administration of cariporide [86] in a dose-dependent manner [87]. Supporting this finding, cariporide prevented endothelial dysfunction in diabetic rats [23]. Furthermore, chronic treatment with cariporide improved endothelial function in obese mice [88]. This evidence suggests that inhibition of NHE-1 could be beneficial in diabetes.

In insulin-resistant animal models, cariporide reduced fasting insulin levels and insulin response to meal tolerance tests indicating improved insulin sensitivity [88]. Furthermore, insulin sensitivity was improved in patients with type 2 diabetes after oral calcium supplementation and was accompanied by decreased activity of NHE-1 in platelets [89]. However, the effect of NHE-1 on insulin sensitivity needs to be investigated.

Improvement of hyperinsulinemia with troglitazone for 4 weeks reduces NHE-1 activity in platelets obtained from fructose-fed borderline hypertensive rats. Moreover, a positive correlation between plasma insulin concentrations and NHE-1 activity ($r = 0.27$, $p = 0.01$) was reported in this study [90].

Insulin inhibited NHE-1 activity in mouse mesenteric artery smooth muscle and endothelial cells through the involvement of H₂O₂ [91]. In this study, the insulin

concentration used (2.5 – 800 nmol/l) was about 10 – 2500 times more than the insulin concentrations encountered in insulin-resistant or diabetic humans (about 0.3 nmol/l). Other studies showed that insulin activates NHE-1 in human monocytes [75] and RBCs [81]. In contrast, others have shown [92,93] that NHE activity is decreased in papillary muscles from the hearts of diabetic rats. This may be due to diabetes-related alterations in Ca/calmodulin protein kinase II. The effect of insulin on NHE-1 activity has raised a controversy whether it is stimulatory or inhibitory. Vaughan-Jones and Swietach in a recent discussion of the matter have concluded that possibly there is a balance between stimulatory and inhibitory actions with a modest overall stimulatory effect [94].

In conclusion, there is evidence that high concentrations of glucose or insulin could activate NHE-1 suggesting that NHE-1 activity might be increased in diabetes. Attenuation of hyperinsulinemia reduces NHE-1 activity. NHE-1 inhibition is also associated with improved insulin sensitivity and prevents endothelial dysfunction. Thus, NHE-1 inhibition in diabetes might have a dual benefit: prevention/treatment of atherogenesis and improved insulin sensitivity. But, this suggestion needs to be proven. It should be noted that in the diabetic myocardium, NHE-1 activity is decreased though it is not clear whether this is a secondary effect.

3.3 Hypertension

Animal and *ex vivo* studies with human peripheral blood cells have documented an increased activity of NHE-1 in primary hypertension [95]. However, genetic linkage studies have excluded the NHE-1 locus from primary hypertension candidate genes [96,97]. Nevertheless, elevated NHE-1 activity has been persistently demonstrated in humans with hypertension and in animals models [98] and it has been suggested that regulation of NHE-1 activity may be affected in hypertension [98]. Alternatively, several newly discovered isoforms of NHE may be responsible for the elevated activity [99].

Inhibition of NHE-1 in spontaneous hypertensive rats with EIPA or amiloride prevented acetylcholine-induced pulmonary artery contraction. However, EIPA or amiloride did not have any effect on the wild-type rats [100]. This finding indicates a different response of hypertensive animals to NHE-1 inhibitors possibly due to altered activity of NHE-1 in these models. So, NHE-1 activity might contribute to the development of hypertension.

NHE overactivity is reported in human lymphocytes obtained from hypertensive patients with obesity or type 2 diabetes [101]. Hyperinsulinemia attenuation following treatment with troglitazone for 4 weeks reduces NHE-1 activity in platelets obtained from fructose-fed borderline hypertensive rats [90].

NHE-1 seems to be associated with hypertension but it still needs to be proven whether NHE-1 contributes to hypertension development or whether NHE-1 inhibition could be used in treatment of hypertension. The discovery

of new NHE isoforms that may be linked with hypertension is exciting and also requires further investigation. Sodium lithium countertransport is elevated in hypertension. A report [102] suggested that a splice variant of NHE1 is responsible for this. However, further studies are necessary to determine the role this splice variant plays in hypertension.

3.4 Dyslipidemia

Several studies have reported that NHE-1 activity is influenced by dyslipidemia. There were reports that in hypercholesterolemic ApoE^{-/-} mice, ischemia and reperfusion caused a smaller infarct size compared with the healthy wild-type mice. This paradoxical finding was explained by the downregulation of NHE-1 protein in cardiomyocytes. Furthermore, histopathology showed no coronary atherosclerosis in these hearts [103]. In accordance with these findings, thrombin-induced NHE-1 activation was reduced in platelets obtained from hereditary hypertensive hypercholesterolemic rats compared with healthy controls [104]. However, there was no difference between these animal models and the healthy controls in basal platelet pHi [104].

It is obvious that high cholesterol levels can alter NHE-1 activity and several studies have proposed underlying mechanisms. LDL inhibits activation of NHE-1 in human platelets in a dose-dependent manner [66], probably through activation of p38^{MAP} kinase [105]. Normally, activation of platelets is accompanied by an initial decrease of pHi, which is counteracted by NHE-1 activity [60]. This mechanism is less pronounced in platelets obtained from humans with familial hypercholesterolemia compared with those obtained from healthy controls [66]. Furthermore, lowering circulating LDL-cholesterol (LDL-C) levels in familial hypercholesterolemia (achieved by LDL apheresis) increased NHE activity [66]. The effect of LDL on NHE-1 resembles the action of NHE-1 inhibitor HOE 694 [66].

High-density lipoprotein (HDL) exerts an opposite effect to LDL on NHE-1 activity [106]. This is an interesting finding as HDL commonly functions in opposition to LDL [107]. HDL₃ enhanced NHE-1 activity after stimulation with acidification even though it had no effect under basal conditions [106]. The proposed mechanism of HDL₃ enhancing effect on NHE-1 is via binding to glycoproteins IIb/IIIa and activation of PKC and phosphatidylcholine-specific phospholipase C [106].

As mentioned earlier, lowering LDL-C enhances NHE-1 activity. However, improvement of hypertriglyceridemia with bezafibrate for 4 weeks did not affect NHE-1 activity [90].

In conclusion, NHE-1 activity seems to be reduced in hypercholesterolemic animal models probably due to increased LDL-C, which reduces NHE-1 activity, or decreased HDL, which enhances NHE-1 activity. However, the clinical relevance of this finding is not yet clear. Biochemically, it has been shown that cholesterol affects NHE-1 activity. Cholesterol

depletion stimulated NHE-1 activity and was associated with relocation outside microdomains [108].

3.5 Smoking

Surprisingly, we could not find any relevant literature linking smoking and NHE-1 activity. However, as NHE-1 activity is influenced by free radicals [109] and cigarette smoking is strongly associated with free radical production [110], an association between NHE-1 activation and smoking cannot be excluded.

3.6 Inflammation

Some emerging risk factors for atherosclerosis have been demonstrated to improve with NHE-1 inhibition. In animal models with heart failure after myocardial infarction (MI), cariporide administration decreased the levels of CRP (C-reactive protein) compared with the placebo group [111]. Inhibition of NHE by either removing extra-platelet Na⁺ or adding an amiloride derivative also decreased arachidonic acid release from platelets [112].

More direct anti-inflammatory effects have also been shown. The anti-inflammatory effect of NHE-1 has been shown *in vivo* and *in vitro* in an animal model; NHE-1 inhibition decreased prostaglandin E2 production through the inhibition of arachidonic acid release and increased the levels of COX-2 protein [113]. In experimental colitis, NHE-1 inhibition resulted in reduction of certain inflammatory markers (myeloperoxidase and IL-1 β) [114]. Further, in endotoxin-exposed endothelial cells, NHE-1 inhibition decreased the inflammatory response of the cells by suppression of IL-8 production, inhibitory- κ B and NF- κ B DNA binding [30]. The anti-inflammatory effects of NHE-1 inhibition and improving CRP levels can result in lower risk of vascular events [115,116]. We showed that ROS trigger increased NHE-1 gene and protein expression [117]. Given the role of ROS in atherosclerosis [118,119], this action may also be relevant.

4. Clinical trials and potential therapeutic applications

The use of several NHE-1 inhibitors in cardiovascular disease has been investigated during the past decade. Animal post-MI studies have shown that NHE-1 inhibition could limit reperfusion injury in the heart [120] (reviewed in [121]) and improve contractile function [122]. Cariporide in combination with metoprolol decreased MI size in animal models [123]. Furthermore, NHE-1 also has a prohypertrophic effect and its inhibition can reduce or prevent heart hypertrophy [121].

A number of clinical studies have examined the efficacy of several NHE-1 inhibitors in the treatment of cardiovascular disease. In the Guard During Ischemia Against Necrosis (GUARDIAN) trial, cariporide was evaluated in patients at risk of myocardial necrosis [124]. Patients with non-ST elevation acute coronary syndrome were enrolled and

randomized to cariporide 20, 80 or 120 mg intravenously 3 times per day or placebo for 2 – 7 days. There was no benefit in the ventricular wall motion score index or in the number of abnormal chords [9]. Analysis of a patients cohort undergoing coronary artery bypass graft surgery (CABG) showed that only the highest dose of cariporide (120 mg) decreased all-cause mortality and MI in the first 36 h postoperatively ($p = 0.005$); the beneficial effect remained for 6 months ($p = 0.033$) [125]. In this cohort, patients were randomized to placebo ($n = 743$) or cariporide 20 mg ($n = 736$), 80 mg ($n = 705$) or 120 mg ($n = 734$). A 1 h intravenous infusion was initiated shortly before surgery and administered every 8 h for 2 – 7 days. The lower doses of cariporide (i.e., 20 or 80 mg) showed no benefit versus placebo [125]. Cariporide seemed to be well tolerated with mild and transient side effects [126].

The EXPEDITION trial (Na⁺/H⁺ Exchange inhibition to Prevent coronary Events in acute cardiac condition) aimed to investigate the effect of cariporide (180 mg in a 1 h pre-CABG loading intravenous dose, then 40 mg/h over 24 h and 20 mg/h over the subsequent 24 h) or placebo on death and MI prevention in patients undergoing CABG ($n = 5761$). There was a decrease in MIs (18.9% placebo group vs 14.4% cariporide group, $p < 0.001$) but an increase in mortality (1.5% placebo group vs 2.2% cariporide group, $p = 0.02$) mainly due to cerebrovascular events [7]. There was an excess of ischemic strokes in this study in the cariporide group (124/2870 vs 67/2839; $p < 0.0001$ calculated by us using the chi-square test with Yates correction). It is of interest that an experimental model has shown that inhibition of NHE-1 activity (using 5-*N,N*-hexamethyleneamiloride or FR183998) prevented the acetylcholine-induced vasodilatation of basilar artery [18].

Another study ($n = 824$) with an inhibitor of NHE-1, zoniopride, showed no benefit in reducing composite cardiovascular end point in patients with multiple risk factors for coronary disease undergoing non-cardiac surgery. In this study, all four groups (placebo, zoniopride 3-, 6-, or 12-mg/kg/day) received a loading dose 1 h before the operation and postoperatively an intravenous maintenance dose for a period of 24 h up to 7 days [127].

In the Evaluation of the Safety and Cardioprotective effects of Eniporide in Acute Myocardial Infarction (ESCAMI) trial, eniporide was infused 10 min before the procedure in patients with ST elevation MI undergoing thrombolytic therapy or angioplasty surgery. Patients were randomized in stage 1 ($n = 430$) to 50, 100, 150 or 200 mg eniporide or placebo and in stage 2 ($n = 959$) to 100 and 150 mg eniporide or placebo. Eniporide versus placebo did not limit MI size or improve clinical outcome [10]. More recently, the pharmacokinetics of eniporide and its metabolite (EMD 112 843) were investigated in healthy subjects and in patients undergoing myocardial reperfusion therapy [128].

The majority of the aforementioned trials have not shown a clinical benefit for NHE-1 inhibitors in patients with

coronary heart disease. A benefit has only been shown in patients undergoing CABG receiving 120 mg dose of cariporide 1 h before CABG and postoperatively every 8 h for 2 – 7 days (GUARDIAN trial) [124]. However, overall mortality was significantly increased ($p = 0.02$) in a later study that also enrolled patients undergoing CABG (EXPEDITION trial) [7]. The opposing findings of these trials could be explained by the differences in the dose administration protocol. Furthermore, the number of patients involved in the studies differed considerably ($n = 1477$ at 120 mg dosage of cariporide in the positive trial (GUARDIAN) and $n = 5761$ in the negative trial (EXPEDITION)).

Newer agents are being investigated as potent inhibitors of NHE-1 with better cardioprotective effects in animal models than the older inhibitors [61,129-133]. Furthermore, an orally administered agent (SMP-300) has also been used in experimental angina and shown better results in infarct size compared with propranolol and nifedipine [134]. Future studies may establish if NHE-1 inhibitors are of clinical benefit. It should be highlighted that all the above-mentioned drugs do not solely inhibit NHE-1 but also other isoforms of NHE with different though lower efficacy [24]. So, it is possible that some of the observed effects in these clinical trials could be caused by inhibition of other NHE isoforms. Also, the presence or absence of bicarbonate in the incubation medium in experiments may influence the results with bicarbonate dependent transport being more of a contributory factor when substrate is available [135].

5. Modulation of NHE-1 activity by clinically used compounds

Several widely used drugs seem to modulate NHE-1 activity. In hypertensive animal models, the administration of enalapril (20 mg/kg/day), nifedipine (10 mg/kg/day) or losartan (40 mg/kg/day) resulted in normalized NHE-1 activity [136]. In young adults with mild hypertension, lisinopril was used for 12 weeks and decreased NHE activity in RBCs [137]. Another ACE inhibitor, quinapril decreased NHE activity in lymphocytes from hypertensive patients after 6 months of treatment [138]. Similarly, enalapril treatment of spontaneous hypertensive rats normalized NHE activity in myocytes [139]. Ramipril and valsartan decreased NHE-1 activity in post-MI animal models mainly due to inhibition of mRNA and protein levels [140]. Long-term (7 days) cilaxapril treatment of hypertensive

animal models prevented the increase in NHE-1 expression [141]. Different results were reported with captopril which increased NHE activity in RBCs from hypertensive patients [142].

In other studies, skeletal muscle NHE was activated by isoprenaline, which stimulates β_2 -adrenoceptors, *in vivo*. This effect was attenuated by nifedipine [143]. Conversely, sildenafil was shown to inhibit cardiac NHE-1 activity and this effect was associated with protecting the myocardium against post-MI remodeling and dysfunction [144].

6. Expert opinion

NHE-1 is involved in the function of all cell types (endothelial cells, monocytes, SMCs and platelets) associated with atherogenesis and with several vascular risk factors (obesity, diabetes, hypertension, dyslipidemia and inflammation). There is evidence that NHE-1 contributes to atherogenesis but it does require further investigation. It is of interest that commonly used drugs (e.g., anti-hypertensive) modulate NHE-1 activity but the clinical relevance of such an effect remains unclear. Furthermore, in many experimental settings, inhibition of NHE-1 has an effect only in states of increased NHE-1 activity. In conclusion, NHE-1 inhibition could contribute to atherogenesis prevention and treatment as long as safe NHE-1 inhibitors are found. Yet, the main limitation of this class of drugs is its lack of specificity.

Acknowledgements

This paper is part of the 03ED29 research project implemented within the framework of the 'Reinforcement Programme of Human Research Manpower' (PENED) and co-financed by National and European Community Funds (25% from the Greek Ministry of Development - General Secretariat of Research and Technology and 75% from the EU - European Social Fund).

Declaration of interest

M Sarigianni is supported by a grant awarded by the Hellenic Atherosclerosis Society. A Tsapas is a Senior Research Fellow of the Tseu Medical Institute within Harris Manchester College, Oxford, United Kingdom, and acknowledges funding support.

Bibliography

1. Putney LK, Denker SP, Barber DL. The changing face of the Na⁺/H⁺ exchanger, NHE1: structure, regulation, and cellular actions. *Annu Rev Pharmacol Toxicol* 2002;42:527-52
2. Fliegel L. The Na⁺/H⁺ exchanger isoform 1. *Int J Biochem Cell Biol* 2005;37:33-7
3. Orłowski J, Grinstein S. Diversity of the mammalian sodium/proton exchanger SLC9 gene family. *Pflugers Arch* 2004;447:549-65
4. Lamon BD, Hajjar DP. Inflammation at the molecular interface of atherogenesis: an anthropological journey. *Am J Pathol* 2008;173:1253-64
5. Libby P, Ridker PM, Hansson GK. Inflammation in atherosclerosis: from pathophysiology to practice. *J Am Coll Cardiol* 2009;54:2129-38
6. Insull W Jr. The pathology of atherosclerosis: plaque development and plaque responses to medical treatment. *Am J Med* 2009;122:S3-14
7. Mentzer RM Jr, Bartels C, Bolli R, et al. Sodium-hydrogen exchange inhibition by cariporide to reduce the risk of ischemic cardiac events in patients undergoing coronary artery bypass grafting: results of the EXPEDITION study. *Ann Thorac Surg* 2008;85:1261-70
8. Teiwes J, Toto RD. Epithelial sodium channel inhibition in cardiovascular disease. A potential role for amiloride. *Am J Hypertens* 2007;20:109-17
9. Tardif JC, Mercier LA, Theroux P. Effect of inhibition of the Na⁺/H⁺ exchanger with cariporide on left ventricular function in acute coronary syndromes: results from the echocardiographic substudy of the GUARDIAN trial. *Can J Cardiol* 2004;20:317-22
10. Zeymer U, Suryapranata H, Monassier JP, et al. The Na⁺/H⁺ exchange inhibitor eniporide as an adjunct to early reperfusion therapy for acute myocardial infarction. Results of the evaluation of the safety and cardioprotective effects of eniporide in acute myocardial infarction (ESCAMI) trial. *J Am Coll Cardiol* 2001;38:1644-50
11. Marcus AJ, Broekman MJ, Drosopoulos JH, et al. Role of CD39 (NTPDase-1) in thromboregulation, cerebroprotection, and cardioprotection. *Semin Thromb Hemost* 2005;31:234-46
12. Yamamoto S, Matsui K, Itoh N, et al. The effect of an Na⁺/H⁺ exchange inhibitor, SM-20550, on ischemia/reperfusion-induced endothelial dysfunction in isolated perfused rat hearts. *Int J Tissue React* 2001;23:1-7
13. Symons JD, Schaefer S. Na⁺/H⁺ exchange subtype 1 inhibition reduces endothelial dysfunction in vessels from stunned myocardium. *Am J Physiol Heart Circ Physiol* 2001;281:H1575-82
14. Muraki S, Morris CD, Budde JM, et al. Blood cardioplegia supplementation with the sodium-hydrogen ion exchange inhibitor cariporide to attenuate infarct size and coronary artery endothelial dysfunction after severe regional ischemia in a canine model. *J Thorac Cardiovasc Surg* 2003;125:155-64
15. Besse S, Tanguy S, Boucher F, et al. Protection of endothelial-derived vasorelaxation with cariporide, a sodium-proton exchanger inhibitor, after prolonged hypoxia and hypoxia-reoxygenation: effect of age. *Eur J Pharmacol* 2006;531:187-93
16. Zatta AJ, Matherne GP, Headrick JP. Adenosine receptor-mediated coronary vascular protection in post-ischemic mouse heart. *Life Sci* 2006;78:2426-37
17. Gumina RJ, Moore J, Schelling P, et al. Na⁺/H⁺ exchange inhibition prevents endothelial dysfunction after I/R injury. *Am J Physiol Heart Circ Physiol* 2001;281:H1260-6
18. Kitazono T, Kamouchi M, Ago T, et al. Role of Na⁺/H⁺ exchanger in dilator responses of rat basilar artery in vivo. *Brain Res* 2001;906:101-6
19. Alnaeb ME, Thompson CS, Seifalian AM, et al. Regional differences in the expression of nitric oxide synthase and specific receptors in the vascular tissues of control and diabetic rabbits: a pilot study. *In Vivo* 2007;21:1069-74
20. Aldakkak M, Stowe DF, Heisner JS, et al. Enhanced Na⁺/H⁺ exchange during ischemia and reperfusion impairs mitochondrial bioenergetics and myocardial function. *J Cardiovasc Pharmacol* 2008;52:236-44
21. Konstantinidis D, Paletas K, Koliakos G, et al. Signaling components involved in leptin-induced amplification of the atherosclerosis-related properties of human monocytes. *J Vasc Res* 2009;46:199-208
22. Xue J, Zhou D, Yao H, et al. Novel functional interaction between Na⁺/H⁺ exchanger 1 and tyrosine phosphatase SHP-2. *Am J Physiol Regul Integr Comp Physiol* 2007;292:R2406-16
23. Wang S, Peng Q, Zhang J, et al. Na⁺/H⁺ exchanger is required for hyperglycaemia-induced endothelial dysfunction via calcium-dependent calpain. *Cardiovasc Res* 2008;80:255-62
24. Masereel B, Pochet L, Laeckmann D. An overview of inhibitors of Na⁺/H⁺ exchanger. *Eur J Med Chem* 2003;38:547-54
25. Vial G, Dubouchaud H, Couturier K, et al. Na⁺/H⁺ exchange inhibition with cariporide prevents alterations of coronary endothelial function in streptozotocin-induced diabetes. *Mol Cell Biochem* 2008;310:93-102
26. Ando K, Takahashi K, Ono A, et al. Possible role of sodium-hydrogen antiport in acetylcholine-induced relaxation of rat aorta. *Biochem Biophys Res Commun* 1991;177:407-13
27. Shankar R, de la Motte CA, DiCorleto PE. Thrombin stimulates PDGF production and monocyte adhesion through distinct intracellular pathways in human endothelial cells. *Am J Physiol* 1992;262:C199-206
28. Buerke U, Pruefer D, Carter JM, et al. Sodium/hydrogen exchange inhibition with cariporide reduces leukocyte adhesion via P-selectin suppression during inflammation. *Br J Pharmacol* 2008;153:1678-85
29. Wang SX, Sun XY, Zhang XH, et al. Cariporide inhibits high glucose-mediated adhesion of monocyte-endothelial cell and expression of intercellular adhesion molecule-1. *Life Sci* 2006;79:1399-404
30. Nemeth ZH, Deitch EA, Lu Q, et al. NHE blockade inhibits chemokine production and NF-kappaB activation in immunostimulated endothelial cells. *Am J Physiol Cell Physiol* 2002;283:C396-403

31. Huck V, Niemeyer A, Goerge T, et al. Delay of acute intracellular pH recovery after acidosis decreases endothelial cell activation. *J Cell Physiol* 2007;211:399-409
32. Park KS, Poburko D, Wollheim CB, et al. Amiloride derivatives induce apoptosis by depleting ER Ca(2+) stores in vascular endothelial cells. *Br J Pharmacol* 2009;156:1296-304
33. Charo IF, Ransohoff RM. The many roles of chemokines and chemokine receptors in inflammation. *N Engl J Med* 2006;354:610-21
34. Viola A, Luster AD. Chemokines and their receptors: drug targets in immunity and inflammation. *Annu Rev Pharmacol Toxicol* 2008;48:171-97
35. Rao GN, de Roux N, Sardet C, et al. Na⁺/H⁺ antiporter gene expression during monocytic differentiation of HL60 cells. *J Biol Chem* 1991;266:13485-8
36. Lei L, Dou L, Yan L, et al. Effect of NHE1 on stem cell differentiation into cardiomyocytes. *Sheng Wu Gong Cheng Xue Bao* 2008;24:1790-5
37. Li X, Karki P, Lei L, et al. Na⁺/H⁺ exchanger isoform 1 facilitates cardiomyocyte embryonic stem cell differentiation. *Am J Physiol Heart Circ Physiol* 2009;296:H159-70
38. Koliakos G, Befani C, Paletas K, et al. Effect of endothelin on sodium/hydrogen exchanger activity of human monocytes and atherosclerosis-related functions. *Ann N Y Acad Sci* 2007;1095:274-91
39. Kaloyianni M, Zolota Z, Paletas K, et al. Cariporide counteracts atherosclerosis-related functions in monocytes from obese and normal individuals. *Obes Res* 2005;13:1588-95
40. Koliakos G, Zolota Z, Paletas K, et al. High glucose concentrations stimulate human monocyte sodium/hydrogen exchanger activity and modulate atherosclerosis-related functions. *Pflugers Arch* 2004;449:298-306
41. Sarigianni M, Bekiari E, Tsapas A, et al. Effect of epinephrine and insulin resistance on human monocytes obtained from lean and obese healthy subjects: a pilot study. *Angiology* 2010; In press
42. Sarigianni M, Bekiari E, Tsapas A, et al. Effect of leptin and insulin resistance on properties of human monocytes in lean and obese healthy subjects. *Angiology* 2010; In press
43. Li Z, Song T, Liu GZ, et al. Inhibitory effects of cariporide on LPC-induced expression of ICAM-1 and adhesion of monocytes to smooth muscle cells in vitro. *Acta Pharmacol Sin* 2006;27:1326-32
44. Schwartz SM, Virmani R, Rosenfeld ME. The good smooth muscle cells in atherosclerosis. *Curr Atheroscler Rep* 2000;2:422-9
45. Bentzon JF, Weile C, Sondergaard CS, et al. Smooth muscle cells in atherosclerosis originate from the local vessel wall and not circulating progenitor cells in apoE knockout mice. *Arterioscler Thromb Vasc Biol* 2006;26:2696-702
46. Doran AC, Meller N, McNamara CA. Role of smooth muscle cells in the initiation and early progression of atherosclerosis. *Arterioscler Thromb Vasc Biol* 2008;28:812-19
47. Boedtker E, Praetorius J, Aalkjaer C. Nbcn1 (SLC4A7) mediates the Na⁺-dependent bicarbonate transport important for regulation of intracellular pH in mouse vascular smooth muscle cells. *Circ Res* 2006;98:515-23
48. Apkon M, Weed RA, Boron WF. Motor responses of cultured rat cerebral vascular smooth muscle cells to intra- and extracellular pH changes. *Am J Physiol* 1997;273:H434-45
49. Aalkjaer C, Mulvany MJ. Effect of changes in intracellular pH on the contractility of rat resistance vessels. *Prog Biochem Pharmacol* 1988;23:150-8
50. Incerpi S, D'Arezzo S, Marino M, et al. Short-term activation by low 17beta-estradiol concentrations of the Na⁺/H⁺ exchanger in rat aortic smooth muscle cells: Physiopathological implications. *Endocrinology* 2003;144:4315-24
51. Berk BC, Vallega G, Muslin AJ, et al. Spontaneously hypertensive rat vascular smooth muscle cells in culture exhibit increased growth and Na⁺/H⁺ exchange. *J Clin Invest* 1989;83:822-9
52. Bkaily G, Nader M, Avedanian L, et al. G-protein-coupled receptors, channels, and Na⁺/H⁺ exchanger in nuclear membranes of heart, hepatic, vascular endothelial, and smooth muscle cells. *Can J Physiol Pharmacol* 2006;84:431-41
53. Wu D, Doods H, Stassen JM. Inhibition of human pulmonary artery smooth muscle cell proliferation and migration by sabiporide, a new specific NHE-1 inhibitor. *J Cardiovasc Pharmacol* 2006;48:34-40
54. Takewaki S, Kuro-o M, Hiroi Y, et al. Activation of Na⁽⁺⁾-H⁺ antiporter (NHE-1) gene expression during growth, hypertrophy and proliferation of the rabbit cardiovascular system. *J Mol Cell Cardiol* 1995;27:729-42
55. Chen YX, O'Brien ER. Ethyl isopropyl amiloride inhibits smooth muscle cell proliferation and migration by inducing apoptosis and antagonizing urokinase plasminogen activator activity. *Can J Physiol Pharmacol* 2003;81:730-9
56. Rich IN, Worthington-White D, Garden OA, et al. Apoptosis of leukemic cells accompanies reduction in intracellular pH after targeted inhibition of the Na⁽⁺⁾/H⁽⁺⁾ exchanger. *Blood* 2000;95:1427-34
57. Schneider D, Gerhardt E, Bock J, et al. Intracellular acidification by inhibition of the Na⁺/H⁺-exchanger leads to caspase-independent death of cerebellar granule neurons resembling paraptosis. *Cell Death Differ* 2004;11:760-70
58. Langer HF, Gawaz M. Platelet-vessel wall interactions in atherosclerotic disease. *Thromb Haemost* 2008;99:480-6
59. Tsiara S, Elisaf M, Jagroop IA, et al. Platelets as predictors of vascular risk: is there a practical index of platelet activity? *Clin Appl Thromb Hemost* 2003;9:177-90
60. Siffert W. Regulation of platelet function by sodium-hydrogen exchange. *Cardiovasc Res* 1995;29:160-6
61. Xu WT, Jin N, Xu J, et al. Design, synthesis and biological evaluation of novel substituted benzoylguanidine derivatives as potent Na⁺/H⁺ exchanger inhibitors. *Bioorg Med Chem Lett* 2009;19:3283-7
62. Jagroop IA, Clatworthy I, Lewin J, et al. Shape change in human platelets: measurement with a channelyzer and visualisation by electron microscopy. *Platelets* 2000;11:28-32
63. Flaumenhaft R. Molecular basis of platelet granule secretion.

- Arterioscler Thromb Vasc Biol 2003;23:1152-60
64. Stelmach H, Rusak T, Tomasiak M. The involvement of the Na⁺/H⁺ exchanger in the formation of microvesicles by porcine platelets. *Haematologia (Budap)* 2002;32:239-52
65. Tomasiak M, Ciborowski M, Stelmach H. The role of Na⁺/H⁺ exchanger in serotonin secretion from porcine blood platelets. *Acta Biochim Pol* 2005;52:811-22
66. Nofer JR, Tepel M, Kehrel B, et al. Low-density lipoproteins inhibit the Na⁺/H⁺ antiport in human platelets. A novel mechanism enhancing platelet activity in hypercholesterolemia. *Circulation* 1997;95:1370-7
67. Barradas MA, Gill DS, Fonseca VA, et al. Intraplatelet serotonin in patients with diabetes mellitus and peripheral vascular disease. *Eur J Clin Invest* 1988;18:399-404
68. Barradas MA, Mikhailidis DP. Serotonin, histamine and platelets in vascular disease with special reference to peripheral vascular disease. *Braz J Med Biol Res* 1992;25:1063-76
69. Bucki R, Pastore JJ, Giraud F, et al. Involvement of the Na⁺/H⁺ exchanger in membrane phosphatidylserine exposure during human platelet activation. *Biochim Biophys Acta* 2006;1761:195-204
70. Tomasiak M, Stelmach H, Rusak T, et al. Vasopressin acts on platelets to generate procoagulant activity. *Blood Coagul Fibrinolysis* 2008;19:615-24
71. Lee KS, Jin YR, Lee JJ, et al. Anti-platelet activity of KR-32560, a novel sodium/hydrogen exchanger-1 inhibitor. *Pharmacol Res* 2006;53:265-70
72. Tomasiak MM, Stelmach H, Bodzenta-Lukaszyk A, et al. Involvement of Na⁺/H⁺ exchanger in desmopressin-induced platelet procoagulant response. *Acta Biochim Pol* 2004;51:773-88
73. Klinkhardt U, Kuczka K, Harder S. Effects of the NHE-1 inhibitor cariporide alone or together with the P2Y₁₂ antagonist AR-C 69331 mx on CD62P expression and formation of platelet-leukocyte aggregates. *Thromb Res* 2003;111:251-7
74. Delva P, Pastori C, Provoli E, et al. Erythrocyte Na⁺-H⁺ exchange activity in essential hypertensive and obese patients: Role of excess body weight. *J Hypertens* 1993;11:823-30
75. Sarigianni M, Bekiari E, Tsapas A, et al. Effect of glucose and insulin on oxidized low density lipoprotein phagocytosis by human monocytes: a pilot study. *Angiology* 2010; In press
76. Paraskevas KI, Liapis CD, Mikhailidis DP. Leptin: a promising therapeutic target with pleiotropic action besides body weight regulation. *Curr Drug Targets* 2006;7:761-71
77. Ravussin E, Tataranni PA. The role of altered sympathetic nervous system activity in the pathogenesis of obesity. *Proc Nutr Soc* 1996;55:793-802
78. Grassi G, Quarti-Trevano F, Seravalle G, et al. Differential sympathetic activation in muscle and skin neural districts in the metabolic syndrome. *Metabolism* 2009;58:1446-51
79. Alvarez GE, Beske SD, Ballard TP, et al. Sympathetic neural activation in visceral obesity. *Circulation* 2002;106:2533-6
80. Sauvage M, Maziere P, Fathallah H, et al. Insulin stimulates NHE1 activity by sequential activation of phosphatidylinositol 3-kinase and protein kinase C zeta in human erythrocytes. *Eur J Biochem* 2000;267:955-62
81. Kaloyianni M, Bourikas D, Koliakos G. The effect of insulin on Na⁺/H⁺ antiport activity of obese and normal subjects erythrocytes. *Cell Physiol Biochem* 2001;11:253-8
82. Sarigianni M, Tsapas A, Mikhailidis DP, et al. Involvement of signaling molecules on Na⁺/H⁺ exchanger-1 activity in human monocytes. *The open Cardiovasc Med J* 2010; In press
83. Konstantinou-Tegou A, Kaloyianni M, Bourikas D, et al. The effect of leptin on Na⁺-H⁺ antiport (NHE 1) activity of obese and normal subjects erythrocytes. *Mol Cell Endocrinol* 2001;183:11-8
84. Bourikas D, Kaloyianni M, Bougoulia M, et al. Modulation of the Na⁺-H⁺ antiport activity by adrenaline on erythrocytes from normal and obese individuals. *Mol Cell Endocrinol* 2003;205:141-50
85. Kaloyianni M, Tsagias N, Liakos P, et al. Stimulation of Na⁺/H⁺ antiport and pyruvate kinase activities by high glucose concentration in human erythrocytes. *Mol Cells* 2004;17:415-21
86. Wang SX, Xiong XM, Song T, et al. Protective effects of cariporide on endothelial dysfunction induced by high glucose. *Acta Pharmacol Sin* 2005;26:329-33
87. Shuang-Xi W, Li-Ying L, Hu M, et al. Na⁺/H⁺ exchanger inhibitor prevented endothelial dysfunction induced by high glucose. *J Cardiovasc Pharmacol* 2005;45:586-90
88. Russell JC, Proctor SD, Kelly SE, et al. Insulin-sensitizing and cardiovascular effects of the sodium-hydrogen exchange inhibitor, cariporide, in the JCR: LA-cp rat and db/db mouse. *J Cardiovasc Pharmacol* 2005;46:746-53
89. Pikilidou MI, Lasaridis AN, Sarafidis PA, et al. Insulin sensitivity increase after calcium supplementation and change in intraplatelet calcium and sodium-hydrogen exchange in hypertensive patients with type 2 diabetes. *Diabet Med* 2009;26:211-19
90. Fujioka Y, Masai M, Tsuboi S, et al. Troglitazone reduces activity of the Na⁺/H⁺ exchanger in fructose-fed borderline hypertensive rats. *Hypertens Res* 2003;26:111-16
91. Boedtkjer E, Aalkjaer C. Insulin inhibits Na⁺/H⁺ exchange in vascular smooth muscle and endothelial cells in situ: involvement of H₂O₂ and tyrosine phosphatase SHP-2. *Am J Physiol Heart Circ Physiol* 2009;296:H247-55
92. Le Prigent K, Lagadic-Gossmann D, Feuvray D. Modulation by pH₀ and intracellular Ca²⁺ of Na⁺-H⁺ exchange in diabetic rat isolated ventricular myocytes. *Circ Res* 1997;80:253-60
93. Feuvray D. Intracellular pH control mechanisms in the diabetic myocardium. *Diabetes Res Clin Pract* 1996;31(Suppl):S87-92
94. Vaughan-Jones RD, Swietach P. Pushing and pulling the cardiac sodium/hydrogen exchanger. *Circ Res* 2008;103:773-5
95. Orlov SN, Adragna NC, Adarichev VA, et al. Genetic and biochemical determinants of abnormal monovalent ion transport in primary hypertension. *Am J Physiol* 1999;276:C511-36
96. Lifton RP, Hunt SC, Williams RR, et al. Exclusion of the Na⁺-H⁺ antiporter

- as a candidate gene in human essential hypertension. *Hypertension* 1991;17:8-14
97. Bobulescu IA, Di Sole F, Moe OW. Na⁺/H⁺ exchangers: physiology and link to hypertension and organ ischemia. *Curr Opin Nephrol Hypertens* 2005;14:485-94
 98. Frolich O, Karmazyn M. The Na-H exchanger revisited: an update on Na-H exchange regulation and the role of the exchanger in hypertension and cardiac function in health and disease. *Cardiovasc Res* 1997;36:138-48
 99. Xiang M, Feng M, Muend S, et al. A human Na⁺/H⁺ antiporter sharing evolutionary origins with bacterial NHaA may be a candidate gene for essential hypertension. *Proc Natl Acad Sci USA* 2007;104:18677-81
 100. Chau WH, Lee WH, Lau WH, et al. Role of Na⁺/H⁺ exchanger in acetylcholine-mediated pulmonary artery contraction of spontaneously hypertensive rats. *Eur J Pharmacol* 2003;464:177-87
 101. Ghigo D, Alessio P, Burzacca S, et al. Na⁺/H⁺ antiporter activity in peripheral blood lymphocytes of obese and type 2 diabetic patients is increased only in the presence of arterial hypertension. *Eur J Clin Invest* 1994;24:166-72
 102. Zerbini G, Maestroni A, Breviaro D, et al. Alternative splicing of NHE-1 mediates Na-Li countertransport and associates with activity rate. *Diabetes* 2003;52:1511-18
 103. Dworschak M, d'Uscio LV, Breukelmann D, et al. Increased tolerance to hypoxic metabolic inhibition and reoxygenation of cardiomyocytes from apolipoprotein E-deficient mice. *Am J Physiol Heart Circ Physiol* 2005;289:H160-7
 104. Zicha J, Kunes J, David-Dufilho M, et al. Cell calcium handling and intracellular pH regulation in hereditary hypertriglyceridemic rats: reduced platelet response to thrombin stimulation. *Life Sci* 1996;59:803-13
 105. Nofer JR, Noll C, Feuerborn R, et al. Low density lipoproteins inhibit the Na⁺/H⁺ antiport in human platelets via activation of p38MAP kinase. *Biochem Biophys Res Commun* 2006;340:751-7
 106. Nofer JR, Tepel M, Kehrel B, et al. High density lipoproteins enhance the Na⁺/H⁺ antiport in human platelets. *Thromb Haemost* 1996;75:635-41
 107. Navab M, Anantharamaiah GM, Reddy ST, et al. HDL as a biomarker, potential therapeutic target, and therapy. *Diabetes* 2009;58:2711-17
 108. Tekpli X, Huc L, Lacroix J, et al. Regulation of Na⁺/H⁺ exchanger 1 allosteric balance by its localization in cholesterol- and caveolin-rich membrane microdomains. *J Cell Physiol* 2008;216:207-20
 109. Konstantinidis D, Paletas K, Koliakos G, et al. The ambiguous role of the Na⁺/H⁺ exchanger isoform 1 (NHE1) in leptin-induced oxidative stress in human monocytes. *Cell Stress Chaperones* 2009;14:591-601
 110. Babizhayev MA, Yegorov YE. Smoking and health: association between telomere length and factors impacting on human disease, quality of life and life span in a large population-based cohort under the effect of smoking duration. *Fundam Clin Pharmacol* 2010; In press
 111. Rungwerth K, Schindler U, Gerl M, et al. Inhibition of Na⁺/H⁺ exchange by cariporide reduces inflammation and heart failure in rabbits with myocardial infarction. *Br J Pharmacol* 2004;142:1147-54
 112. Sweatt JD, Johnson SL, Cragoe EJ, et al. Inhibitors of Na⁺/H⁺ exchange block stimulus-provoked arachidonic acid release in human platelets. Selective effects on platelet activation by epinephrine, ADP, and lower concentrations of thrombin. *J Biol Chem* 1985;260:12910-19
 113. Kamachi F, Ban HS, Hirasawa N, et al. Inhibition of lipopolysaccharide-induced prostaglandin E2 production and inflammation by the Na⁺/H⁺ exchanger inhibitors. *J Pharmacol Exp Ther* 2007;321:345-52
 114. Khan I, Oriowo MA, Anim JT. Amelioration of experimental colitis by Na-H exchanger-1 inhibitor amiloride is associated with reversal of IL-1ss and ERK mitogen-activated protein kinase. *Scand J Gastroenterol* 2005;40:578-85
 115. Ridker PM, Morrow DA, Rose LM, et al. Relative efficacy of atorvastatin 80 mg and pravastatin 40 mg in achieving the dual goals of low-density lipoprotein cholesterol < 70 mg/dl and C-reactive protein < 2 mg/l: an analysis of the PROVE-IT TIMI-22 trial. *J Am Coll Cardiol* 2005;45:1644-8
 116. Ridker PM, Danielson E, Fonseca FA, et al. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med* 2008;359:2195-207
 117. Akram S, Teong HF, Fliegel L, et al. Reactive oxygen species-mediated regulation of the Na⁺/H⁺ exchanger 1 gene expression connects intracellular redox status with cells' sensitivity to death triggers. *Cell Death Differ* 2006;13:628-41
 118. Harrison D, Griendling KK, Landmesser U, et al. Role of oxidative stress in atherosclerosis. *Am J Cardiol* 2003;91:7A-11A
 119. Bonomini F, Tengattini S, Fabiano A, et al. Atherosclerosis and oxidative stress. *Histol Histopathol* 2008;23:381-90
 120. Imahashi K, Mraiche F, Steenbergen C, et al. Overexpression of the Na⁺/H⁺ exchanger and ischemia-reperfusion injury in the myocardium. *Am J Physiol Heart Circ Physiol* 2007;292:H2237-47
 121. Fliegel L. Regulation of the Na⁽⁺⁾/H⁽⁺⁾ exchanger in the healthy and diseased myocardium. *Expert Opin Ther Targets* 2009;13:55-68
 122. Karmazyn M, Sawyer M, Fliegel L. The Na⁽⁺⁾/H⁽⁺⁾ exchanger: a target for cardiac therapeutic intervention. *Curr Drug Targets Cardiovasc Haematol Disord* 2005;5:323-35
 123. Wang P, Zaragoza C, Holman W. Sodium-hydrogen exchange inhibition and beta-blockade additively decrease infarct size. *Ann Thorac Surg* 2007;83:1121-7
 124. Theroux P, Chaitman BR, Danchin N, et al. Inhibition of the sodium-hydrogen exchanger with cariporide to prevent myocardial infarction in high-risk ischemic situations. Main results of the GUARDIAN trial. Guard during ischemia against necrosis (GUARDIAN) investigators. *Circulation* 2000;102:3032-8
 125. Boyce SW, Bartels C, Bolli R, et al. Impact of sodium-hydrogen exchange inhibition by cariporide on death or myocardial infarction in high-risk CABG surgery patients: results of the CABG surgery cohort of the GUARDIAN study. *J Thorac Cardiovasc Surg* 2003;126:420-7

126. Chaitman BR. A review of the guardian trial results: clinical implications and the significance of elevated perioperative ck-mb on 6-month survival. *J Card Surg* 2003;18(Suppl 1):13-20
127. Fleisher LA, Newman MF, St Aubin LB, et al. Efficacy of zonisipride, an Na/H exchange ion inhibitor, for reducing perioperative cardiovascular events in vascular surgery patients. *J Cardiothorac Vasc Anesth* 2005;19:570-6
128. Bhattaram VA, Nagaraja NV, Peters T, et al. Population pharmacokinetics of eniporide and its metabolite in healthy subjects and patients with acute myocardial infarction. *J Clin Pharmacol* 2005;45:631-9
129. Zhang R, Dong J, Xu YG, et al. Synthesis and bioactivity of substituted indan-1-ylideneaminoguanidine derivatives. *Eur J Med Chem* 2009;44:3771-6
130. Zhang R, Lei L, Xu YG, et al. Benzimidazol-2-yl or benzimidazol-2-ylthiomethyl benzoylguanidines as novel na⁺/h⁺ exchanger inhibitors, synthesis and protection against ischemic-reperfusion injury. *Bioorg Med Chem Lett* 2007;17:2430-3
131. Mao D, Xu Y, Hu X, et al. Synthesis and Na⁽⁺⁾/H⁽⁺⁾ exchanger-1 inhibitory activity of substituted (quinolinecarbonyl) guanidine derivatives. *Chem Biodivers* 2009;6:1727-36
132. Kusumoto K, Igata H, Abe A, et al. In vitro and in vivo pharmacology of a structurally novel Na⁺/H⁺ exchange inhibitor, T-162559. *Br J Pharmacol* 2002;135:1995-2003
133. Fukumoto S, Imamiya E, Kusumoto K, et al. Novel, non-acylguanidine-type Na⁽⁺⁾/H⁽⁺⁾ exchanger inhibitors: synthesis and pharmacology of 5-tetrahydroquinolinylidene aminoguanidine derivatives. *J Med Chem* 2002;45:3009-21
134. Yamamoto S, Matsui K, Sasabe M, et al. Effect of an orally active Na⁺/H⁺ exchange inhibitor, SMP-300, on experimental angina and myocardial infarction models in rats. *J Cardiovasc Pharmacol* 2002;39:234-41
135. Chesler M. Regulation and modulation of pH in the brain. *Physiol Rev* 2003;83:1183-221
136. Alvarez BV, Ennis IL, De Hurtado MC, et al. Effects of antihypertensive therapy on cardiac sodium/hydrogen ion exchanger activity and hypertrophy in spontaneously hypertensive rats. *Can J Cardiol* 2002;18:667-72
137. Falkner B, Canessa M, Anzalone D. Effect of angiotensin converting enzyme inhibitor (lisinopril) on insulin sensitivity and sodium transport in mild hypertension. *Am J Hypertens* 1995;8:454-60
138. Fortuno A, Tisaire J, Lopez R, et al. Angiotensin converting enzyme inhibition corrects Na⁺/H⁺ exchanger overactivity in essential hypertension. *Am J Hypertens* 1997;10:84-93
139. Ennis IL, Alvarez BV, Camilion de Hurtado MC, et al. Enalapril induces regression of cardiac hypertrophy and normalization of pH_i regulatory mechanisms. *Hypertension* 1998;31:961-7
140. Sandmann S, Yu M, Kaschina E, et al. Differential effects of angiotensin AT1 and AT2 receptors on the expression, translation and function of the Na⁺/H⁺ exchanger and Na⁺-HCO₃⁻ symporter in the rat heart after myocardial infarction. *J Am Coll Cardiol* 2001;37:2154-65
141. Khan I, Al-Qattan KK, Alnaqeeb MA, et al. Altered expression of Na⁽⁺⁾/H⁽⁺⁾ exchanger isoforms 1 and 3 in clipped and unclipped kidneys of a 2-kidney-1-clip Goldblatt model of hypertension. *Nephron* 2002;92:346-55
142. Lijnen P, Arabidze CG, Petrov V. Effect of angiotensin-converting enzyme inhibition on Na⁽⁺⁾/H⁽⁺⁾-exchange and Ca⁽²⁺⁾-efflux in human erythrocytes. *Methods Find Exp Clin Pharmacol* 1997;19:125-8
143. Syme PD, Brunotte F, Green Y, et al. The effect of beta 2-adrenoceptor stimulation and blockade of I-type calcium channels on in vivo Na⁺/H⁺ antiporter activity in rat skeletal muscle. *Biochim Biophys Acta* 1991;1093:234-40
144. Perez NG, Piaggio MR, Ennis IL, et al. Phosphodiesterase 5a inhibition induces Na⁺/H⁺ exchanger blockade and protection against myocardial infarction. *Hypertension* 2007;49:1095-103

Affiliation

Maria Sarigianni^{1,2}, Apostolos Tsapas^{1,3}, Dimitri P Mikhailidis², Martha Kaloyianni⁴, George Koliakos⁵, Larry Fliegel⁶ & Konstantinos Paletas^{†7}

[†]Author for correspondence
¹Aristotle University of Thessaloniki, Medical School, Metabolic Diseases Unit, Second Department of Internal Medicine, Greece
²University College London (UCL), University College London Medical School, Royal Free Hospital campus, Department of Clinical Biochemistry (Vascular Disease Prevention Clinics), London NW3 2QG, UK
³University of Oxford, The Tseu Medical Institute, Harris Manchester College, Oxford, UK
⁴Aristotle University of Thessaloniki, School of Biology, Laboratory of Animal Physiology, Department of Zoology, Greece
⁵Aristotle University of Thessaloniki, Medical School, Department of Biological Chemistry, Greece
⁶University of Alberta, Department of Biochemistry, Faculty of Medicine, Edmonton, Alberta T6G 2H7, Canada
⁷Professor, Aristotle University of Thessaloniki, Medical School, Metabolic Diseases Unit, Second Department of Internal Medicine, 15 Ag. Sofias str, 54623, Thessaloniki, Greece
 Tel: +00302310892751;
 Fax: +00302310992937;
 E-mail: paletas@med.auth.gr

Expert Opin. Investig. Drugs Downloaded from informahealthcare.com by 174.3.239.171 on 11/28/10 For personal use only.